

Please replace the paragraph beginning on page 15, line 13, with the following rewritten paragraph:

--**Figures 7A and 7B** show that decreased dietary protein intake correlates with less mitochondrial DNA damage in control and apoE mice. Control and apoE mice were fed either a 16% or 24% protein diet for 4 weeks, commencing at 6 weeks of age, and mitochondrial DNA damage was assessed by QPCR. **Figure 7A** illustrates the relative amplification of control and apoE mitochondrial DNA in aortas relative to the 24% protein fed control group. Less amplification product signifies increased mitochondrial DNA damage. Asterisks (*) indicate a significant difference between the 24% and 16% protein fed mice. Students t' test P values are in parentheses. **Figure 7B** illustrates the relative amplification of control and apoE mitochondrial DNA in the left ventricle relative to the 24% protein fed control group. Less amplification product signifies increased mitochondrial DNA damage. Asterisks (*) indicate a significant difference between the 24% and 16% protein fed mice. Students t' test P values are in parentheses.--

Please replace the paragraph beginning on page 16, line 8, with the following rewritten paragraph:

--**Figure 8** shows that lipid peroxidation is significantly increased in apoE compared to control mice, but lipid peroxidation is significantly increased in control mice fed the western (21% fat) diet. Bar graph illustrates the level of lipid peroxidation in apoE and control mice fed chow (4% fat) or western (21%) diets. Values are expressed relative to the chow fed control mice. While apoE had significantly higher levels of lipid peroxidation compared to control mice regardless of diet, there were

no significant differences in the level of lipid peroxidation among the apoE fed the chow or western diets. By contrast, the control mice fed the western diet had significantly higher levels of lipid peroxidation (indicated by “*”) compared to control mice fed the chow diet.--

Please replace the paragraph beginning on page 16, line 20, with the following rewritten paragraph:

--**Figure 9** shows QPCR of age-matched normal and atherosclerotic aortic mtDNA from humans. The 16.2 kb (upper row) and 0.22 kb (bottom row) products represent the long and short human QPCR amplicons, respectively. Methods for QPCR from human mtDNA have been previously described (Ballinger, 1996, 1999; Yakes, 1997).--

Please replace the paragraph beginning on page 17, line 5, with the following rewritten paragraph:

--**Figures 10A and 10B** show that hybrid apoE (-/-), SOD2 (+/-) mice have increased atherosclerotic lesions and mtDNA damage. In **Figure 10A**, whole aortas from hybrid apoE (-/-), SOD2 (+/-) and apoE (-/-) littermates were oil-red-O stained. The upper left panel shows an oil-red-O stained aorta from a 17 week-old apoE (-/-), SOD2 (+/-) male (4% fat diet), and the upper right panel shows the results from an apoE (-/-) male littermate (4% fat diet). Lower left and right panels show stained aortas from 34 week-old apoE (-/-), SOD2 (+/-) and apoE (-/-) littermates, respectively. Black arrows indicate presence of large atherosclerotic lesion in the apoE (-/-), SOD2 (+/-) and apoE (-/-), (not yet present in the 17 week-old apoE (-/-) littermate), and the red arrow in each panel point to an example of smaller lesion formed at arterial branches, that were more frequent and characteristics of the apoE (-/-), SOD2 (+/-) mice. **Figure**

10B is a bar graph depicting atherosclerotic lesion frequencies from apoE (/), SOD2 (+/-) and apoE (-/-) mice whole aortas, visualized by oil-red-O staining (n=4).--

Please replace the paragraph beginning on page 18, line 1, with the following rewritten paragraph:

--**Figure 11** shows the increased incidence of mitochondrial DNA damage in patients with risk factors for myocardial infarction. Blood samples were taken at the time of cardiac catheterization to measure mitochondrial DNA damage by QPCR. For this figure, values of mitochondrial DNA damage above the mean for a "normal" population were considered increased. The bar graphs show the percentage of patients in each category with increased mitochondrial DNA damage. The occurrence of increased mitochondrial DNA damage is more frequent in subjects with hypertension, cigarette smoking, diabetes and combinations of these risk factors for myocardial infarction.--

Please replace the paragraph beginning on page 18, line 12, with the following rewritten paragraph:

--**Figure 12** shows mitochondrial DNA damage before and after a 20 mile "training run". Blood samples were taken, a buffy coat was obtained and mitochondrial DNA damage was assessed by QPCR. Although mitochondrial DNA damage is present immediately after the training run, this quickly returns to normal values as shown here.--

Please replace the paragraph beginning on page 18, line 18, with the following rewritten paragraph:

mitochondrial DNA damage compared to the age matched control group (1.325 ± 0.257 lesions/10 kb versus 0.453 ± 0.162 lesions/10 kb; $P=0.007$). These same patterns were also observed in tissue of the left ventricle of the heart. (data not shown). For example, the 10 week old apoE mice had a 67% decrease in amplification compared to the zero class control (0.685 ± 0.093 lesions/10 kb versus 0.0 ± 0.048 lesions/10 kb; $P < 0.001$), while the 34 week old apoE mice sustained approximately 4-fold higher estimated mitochondrial DNA lesions compared to the age-matched controls (0.819 ± 0.151 lesions/10 kb versus 0.213 ± 0.295 lesions/10 kb, $P = 0.056$).--

Please replace the paragraph beginning on page 54, line 4, with the following rewritten paragraph:

--The effects of the western diet were also assessed in apoE and control mice. In the control mice, the western diet was associated with higher levels of mitochondrial DNA damage in the 10 week-old control group aortas compared to chow fed control mice (0.31 ± 0.17 lesions/10 kb versus 0.0 ± 0.20 lesions/10 kb, western and chow, respectively), however, this difference was less significant ($P = 0.25$). Such differences were less obvious in the 34 week old control mice (i.e. 0.45 ± 0.16 lesions/10 kb and 0.49 ± 0.25 lesions/10 kb in chow versus western, respectively, $P = 0.90$). Similarly, the 10 week old control mice showed increased estimated mitochondrial DNA damage with the western diet in heart (58% decrease in amplification, 0.54 ± 0.23 lesions/10 kb and 0.0 ± 0.05 lesions/10 kb in western versus chow, respectively), yet similar with the results seen in the aorta, no obvious differences were seen in the 34 week old control mice (0.21 ± 0.29 lesions/10 kb and 0.25 ± 0.21 lesions/kb in chow versus western, respectively). By contrast, no apparent increases in mitochondrial

DNA damage accompanied the western diet in either aorta or heart tissues from apoE mice. Hence, increased dietary fat appeared to be associated with increased mitochondrial DNA damage in the 10 week old control mice only.--

Please replace the paragraph beginning on page 55, line 20, with the following rewritten paragraph:

--Because no significant differences in mitochondrial DNA damage were observed between the chow and western diets with matched genotype and age, the effects of protien in the 10 week old mice were investigated. Both apoE and c57B1control mice were fed either 16% or 24% protein diets for 4 weeks, commencing at 6 weeks of age. While both diets resulted in apoE still having greater levels of mitochondrial DNA damage compared to controls (aorta $P = 0.015$; heart $P = 0.005$), the lower protein diet resulted in a significant decrease in damage in both control ($P = 0.007$) and apoE ($P < 0.001$) in aorta (Figure 7). This same trend was also observed in apoE heart ($P = 0.002$). Hence, the lower protein diet was associated with decreased mitochondrial DNA damage in both types of mice.--

Please replace the paragraph beginning on page 58, line 8, with the following rewritten paragraph:

--Lipid peroxidation levels were measured by determining the levels of MDA and 4-HNE. The hypercholesteremic apoE mice had significantly increased levels of lipid peroxides compared to control mice ($P < 0.05$). However, in contrast with the results observed with the total cholesterol levels, the degree of lipid peroxidation did not significantly increase in apoE mice on the western compared to the chow diet (Figure 8).

By contrast, the control mice fed the western diet sustained increased lipid peroxidation compared to the chow diet ($P < 0.05$). Hence, while the apoE mice always had the highest levels of lipid peroxides compared to the control mice, the degree of lipid peroxidation in apoE did not increase when fed the western diet compared to the chow diet.--

Please replace the paragraph beginning on page 60, line 4, with the following rewritten paragraph:

--QPCR analysis revealed significantly increased mtDNA damage in atherosclerotic tissues compared with healthy control group aortas (Figures 9 and Table 8). By contrast, a marker for nuclear DNA damage, the *β-globin* gene cluster, did not sustain significant damage in atherosclerotic aorta compared with controls ($P = 0.15$). Consequently, the mtDNA sustained significant and preferential damage in human atherosclerotic aortas. These findings were consistent with the notion that mtDNA damage is an integral feature of atherogenesis. Because increased mtDNA damage was a consistent feature in both human and mouse atherogenesis, it was hypothesized that mitochondrial dysfunction is an important event in atherogenesis, possibly modulating the phenotype of vascular cells that are integral to atherosclerotic lesion formation.--

Please replace the paragraph beginning on page 62, line 11, with the following rewritten paragraph:

--For analysis of atherogenesis and mtDNA damage, apoE (-/-), SOD2 (+/-) mice and apoE (-/-) controls were fed a 4% fat diet and sacrificed at 17 weeks of age. Assessment of functional SOD2 activity in hybrid apoE (-/-), SOD2 (+/-) aortas demonstrated a 44% reduction of activity compared with "normal" apoE (-/-) mice (data not shown; $P <$

0.001). Histological examination of atherosclerotic lesion development in these mice revealed that apoE (-/-), SOD2 (+/-) mice developed earlier lesions at physiologically relevant sites, including in the ascending aorta at the origin of the cerebral vessels. However, the most impressive finding was that at 17 weeks of age, oil-red-O positive lesions were present throughout the descending aorta at virtually every branch point of the spinal arteries in the apoE (-/-), SOD2 (+/-) mice (**Figure 10A**). Analogous lesions were rare in both 17-week-old and 34-week-old apoE (-/-) mice. Overall, the apoE (-/-), SOD2 (+/-) mice had 2.5 fold as many atherosclerotic lesions ($P = 0.02$) compared with apoE (-/-) littermates (**Figure 10B**). Although the small size of atherosclerotic lesions at the branch points precluded direct measurement of mtDNA damage at these sites, quantification of aortic mtDNA damage through QPCR revealed increased aortic mtDNA damage in the apoE (-/-), SOD2 (+/-) mice compared with their age-matched apoE (-/-) littermates (**Table 8**; $P = 0.006$). Thus, while providing further evidence that mitochondrial dysfunction resulting from oxidative mtDNA damage is important in atherogenesis, these findings also supports the notion that, under normal circumstances, decreased expression of SOD2 in response to turbulent flow at arterial branch points promotes atherogenesis at these sites.--

Please replace the paragraph beginning on page 77, line 9, with the following rewritten paragraph:

--Patient who were undergoing cardiac catheterization were recruited for an *in vivo* study of mitochondrial DNA damage. All signed informed consent for an additional blood sample to be taken at the time of cardiac catheterization. In all, there were 75 patient. White blood cells were isolated by standard techniques (the "buffy coat" preparation), and DNA